

#### **INTENDED USE**

Lyse-G is a stabilized and microfiltered lysing reagent, for stromatolysis of Erythrocytes (RBCs), for quantitative determination of Leucocytes (WBCs), Leucocyte three-part differentials (LYM, MID, NEUT) and Hemoglobin (HGB) concentration in human blood, using 3 Part Genius Hematology analyzers.

Please refer to the analyzer's User Manual for further information.

#### **ACTIVE INGREDIENTS**

Sodium Chloride: 1.2%, Potassium Chloride: 0.03%, Quaternary Ammonium salt: 3.0%, Preservative: 0.05 %

### **DIRECTIONS FOR USE**

- 1. Personnel installing the reagent, should be a trained laboratory professional.
- 2. Open the container cover, pull the connector of the container out and connect it with the corresponding Lyse tubing as per instructions given in the analyzer's User Manual.
- 3. Avoid contamination of the tubing and the reagent.
- 4. Do not pour and mix the remains of a reagent from a container, into another one.
- 5. Prime thoroughly with the new reagent and measure background values according to the analyzer's User Manual.
- 6. When installing a new lot of the reagent, recalibrate the instrument as specified in the User Manual.

# **PRECAUTIONS**

- a. Lyse G is for in vitro diagnostic use only.
- b. Lyse G is cyanide-free, azide-free and environmental friendly reagent and does not contain harmful ingredients.
- c. Do not inhale the reagent or pipette by mouth.
- d. Avoid contact with skin/eyes. In case of contact, wash off immediately with plenty of water and seek medical attention, as early as possible.
- e. Do not use if package is damaged.
- f. Keep the reagent container closed when not in use.
- g. Wear protective gloves when handling the reagent.
- h. Please refer to the MSDS associated with the reagent.
- i. Use Good Laboratory Practices (GLP) when handling the reagent.

### **SPECIMEN COLLECTION AND STORAGE**

Blood specimens should be collected by vein puncture in EDTA anticoagulant.

Specimens for hematological analysis may be stored at  $15^{\circ} - 30^{\circ}$ C for 2 - 4 hours or at  $2^{\circ} - 8^{\circ}$ C for 24 hours after collection.

### **REAGENT PREPARATION**

This reagent is ready for use and can be connected to the analyzer straight from the container; no special reagent preparation is required.

Phone: +91-11-41727222, +91-11-41084222

Email: info@vanguarddiagnostics.com

Website: www.vanguarddiagnostics.com

# STORAGE, STABILITY AND DISPOSAL

- Store Lyse -G between 15° and 30°C.
- The unopened reagent is stable up to the expiry date mentioned on the label when stored between 15°C 30°C.
- Use Lyse G within 06 months of first opening.
- Do not use reagent beyond the expiry date printed on the container label.
- Discard opened container after 06 months.
- Do not use reagent once frozen.
- Dispose waste product in accordance with local Government regulations.