

#### **INTENDED USE**

Rinse-M is a buffered, stabilized and micro-filtered electrolyte reagent, for the automated cleaning the bath and tubing, using 3 part Mindray Hematology analyzers.

Please refer to the analyzer's User Manual for further information.

#### **ACTIVE INGREDIENTS**

Tris Buffer: 1.0 %, pH 7.5, Sodium Sulphate: 1.0%, Sodium Chloride: 0.5%, Triton X 100: 0.5%, Preservative: 0.10%.

### **DIRECTIONS FOR USE**

- **1.** Personnel installing the reagent, should be a trained laboratory professional.
- 2. Open the container cover, pull the connector of the container out and connect it with the corresponding Rinse tubing as per instructions given in the analyzer's User Manual.
- **3.** Avoid contamination of the tubing and the reagent.
- **4.** Do not pour and mix the remains of a reagent from a container, into another one.
- 5. Prime thoroughly with the new reagent and measure background values according to the analyzer's User Manual.
- 6. When installing a new lot of the reagent, recalibrate the instrument as specified in the User Manual.

### **PRECAUTIONS**

- a. Rinse M, is for *in vitro* diagnostic use only.
- b. Rinse-M is environmental friendly and does not contain harmful ingredients.
- c. Do not inhale the reagent or pipette by mouth.
- d. Avoid contact with skin/eyes. In case of contact, wash off immediately with plenty of water and seek medical attention, as early as possible.
- e. Do not use if package is damaged.
- f. Keep the reagent container closed when not in use.
- g. Wear protective gloves when handling the reagent.
- h. Please refer to the MSDS associated with the reagent.
- i. Use Good Laboratory Practices (GLP) when handling the reagent.

# **SPECIMEN COLLECTION AND STORAGE**

Blood specimens should be collected by vein puncture in EDTA anticoagulant.

Specimens for hematological analysis may be stored at  $15^{\circ} - 30^{\circ}$ C for 2 - 4 hours or at  $2^{\circ} - 8^{\circ}$ C for 24 hours after collection.

## **REAGENT PREPARATION**

This reagent is ready for use and can be connected to the analyzer straight from the container; no special reagent preparation is required.

Phone: +91-11-41727222, +91-11-41084222

Email: info@vanguarddiagnostics.com

Website: <u>www.vanguarddiagnostics.com</u>

## STORAGE, STABILITY AND DISPOSAL

- Store Rinse M between 15° and 30°C.
- The unopened reagent is stable up to the expiry date mentioned on the label when stored between 15°C 30°C.
- Use Rinse M within 06 months of first opening.
- Do not use reagent beyond the expiry date printed on the container label.
- Discard opened container after 06 months.
- Do not use reagent once frozen.
- Dispose waste product in accordance with local Government regulations.